Montana Noxious
Weed Education



Whitetop




Photo Credits: large photo and bottom middle photo courtesy Chris Evans, Illinois Wildlife Action Plan, Bugwood.org; bottom middle photo courtesy of Pedro Tenorio-Lezama, Bugwood.org; bottom right photo courtesy of Julia Scher, USDA APHIS PPQ, Bugwood.org

Whitetop

Genus: Cardaria Species: draba Priority Listing: 2B

Perennial

Whitetop is native to Russia and Eurasia and was accidentally introduced into the United States in contaminated crop seed in the early 1900's. Whitetop was first discovered in Gallatin County, Montana, in 1916. There are actually three different Cardaria species that are collectively referred to as "whitetop": C. pubescens, C. chalapensis, and C. draba.

Whitetop typically inhabits open, sunny areas that have been recently disturbed. These areas include over grazed pastures, waste areas, roadsides, and open grasslands.

Whitetop is a creeping perennial and can grow up to 2' in height with rhizomatous roots, which can extend 2' down and up to 12' out.

The plant is silvery gray-green in color and its leaves are lance to arrowhead shaped and covered with fine hairs. The lower leaves are stalked and the upper leaves clasp the stem. Whitetop produces numerous white flowers that have 4 petals on ½" long stalks. These dense clusters of flowers create a white, flat topped appearance.

Germination: is the process by which plants, fungi, and bacteria emerge from seeds and spores, and begin growth

Inhabits: to live in, reside in, or be present in Integrated Weed Management Techniques: the combination of multiple management tools to reduce a pest population to an acceptable level while preserving the quality of existing habitat, water, and other natural resources

One plant is capable of producing 1,200 to 4,800 seeds. Seeds are produced in fruiting pods that contain 2 chambers, each capable of producing one seed. Seeds are oval in shape and reddish brown in color. Seeds can remain viable in the soil for up to 4 years! Seeds are scattered when the side wall of the seed pod ruptures or when the seed pods fall to the ground. Seeds can germinate in the fall or the spring.

Whitetop is difficult to control because it can reproduce through rhizomes as well as seeds.

Whitetop often requires many years of repetitive herbicide treatments; integrated weed management techniques such as hand pulling and herbicides when combined work best in controlling whitetop.



Herbicide: a chemical substance used to kill or destroy plants, likely to be used on weeds

Perennial: a plant whose life spans several years

Rhizomes: a perennial underground stem that usually grows horizontally

Viable: capable of growing or developing